

DR. LIONEL ROSTAING (Orcid ID : 0000-0002-5130-7286)

Article type : Letter to the Editor

## Why the immune system fails to mount an adaptive immune response to a Covid -19 infection

Ahmadpoor P. <sup>1,2</sup> MD, Rostaing L. <sup>3,4</sup> MD, PhD.

<sup>1</sup> Service de Néphrologie, CHU Carémeau Nîmes, France

<sup>2</sup> Labafi Nejad hospital, Shahid Beheshti medical university, Tehran, Iran

<sup>3</sup> Service de Néphrologie, Hémodialyse, Aphérèses et Transplantation Rénale, CHU Grenoble-Alpes, France

<sup>4</sup> Université Grenoble-Alpes, France

### Address for corresponding author:

Lionel Rostaing, MD, PhD

Service de Néphrologie, Dialyse, Aphérèses et Transplantation Rénale,

CHU Grenoble-Alpes

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/TRI.13611](#)

This article is protected by copyright. All rights reserved

CS 10217

38043 Grenoble Cedex 09, France

Tel: +33 4 76 76 54 60

Fax +33 4 76 76 52 63

Email: lrostaing@chu-grenoble.fr

**Running title:** Immune response and Covid-19 infection

**Key words:** Covid-19; immune response; kidney transplantation; tocilizumab; cytokine storm.

Dear Editors,

The Covid-19 pandemic has already affected many thousands of people and has become the greatest health challenge worldwide [1]. The range of clinical presentations varies from asymptomatic and mild clinical symptoms to acute respiratory-distress syndrome (ARDS) and death. Due to the unknown number of asymptomatic viral-shedding and pauci-symptomatic people in the community, the total number of infections is uncertain. As yet, effective treatment is unavailable. However, the results of preliminary studies and clinical trials are improving our understanding of the pathogenesis and treatment of Covid-19. Herein, we share our personal opinions on the immunopathogenesis of this infection. The virus enters the cell via the angiotensin-converting enzyme-2 (ACE-2) and is sensed (essentially) by Toll-like receptor -7 (TLR -7), which exists in endosomes [2]. TLR -7 activation leads to the production of alpha-interferon, TNF-alpha, and the secretion of interleukin (IL)-12 and IL-6. This results in the formation of CD8<sup>+</sup>-specific cytotoxic T-cells and, through the CD4<sup>+</sup> helper T-cell, leads to the formation of antigen-specific B-cells and antibody production [3,4]. This adaptive immune response controls the viral infection and determines clinical recovery.

It seems that when the body is unable to produce an adequate adaptive response against the virus, the persistent innate-induced inflammation can then lead to a cytokine storm, ARDS, and diffuse organ involvement [5]. Apart from importance of comorbidities like hypertension, diabetes mellitus and coronary heart disease in increasing the risk of mortality, immunosenescence must play a crucial role. With aging, the population of naïve T-cells shrinks while antigen-experienced, memory T-cells comprise an important portion of T-cells population (6,7). It means that the ability of immune system of elderly to respond to previously exposed pathogens is more preserved compared to never-exposed pathogens. In contrast, in children there is huge number of naïve T-cells ready to be educated by new pathogens. This can be one of the explanations for Covid19 milder presentation and significantly lower mortality in children compared to elderly. This situation is close to solid organ transplanted (SOT) patients because our routine immunosuppressive drugs are effective against naïve T-cells but not able to block memory T-/B-cells. During TLR-7 signaling-related cytokine secretion, IL-6 may play an important role. Interleukin-6 has a pleotropic role across the immune system. It is crucial in the formation of follicular helper T-cells, TH17 subset deviation, and for the formation of long-lived plasma-cells. However, IL-6 can block CD8<sup>+</sup> cytotoxic T-cells by inhibiting the secretion of gamma-interferon. Moreover, IL-6 by inducing suppression of cytokine signaling

(SOCS-3) and increasing the expression of PD-1, can paralyze the cell-mediated antiviral response during a cytokine storm [8]. Among the current medications used to manage Covid-19, although results from the antiviral drugs are still inconclusive [9], studies on chloroquine and its less toxic derivative hydroxychloroquine have been associated with shortening viral replication [10]. Their mechanisms of action are to block viral replication in the endosome by preventing endosomal acidification and endolysosomal fusion. Moreover, hydroxychloroquine blocks entry of the virus into cells. In addition, hydroxychloroquine can block TLR7 and TLR9 signaling and, thus, can reconstruct the CD8<sup>+</sup> cytotoxic viral response [11,12]. The addition of azithromycin further reduces the nasopharyngeal Covid-19 virus PCR, as shown by Gautret et al. This is probably through azithromycin immunomodulatory activity via blocking IL-6 and TNF-alpha [13].

Furthermore, in severe Covid-19 cases, e.g., in ARDS, a preliminary report from China showed that the addition of tocilizumab was efficient via blocking IL-6 activity [14]. A clinical trial in Italy (TOCIVID-19) is ongoing (NCT04317092).

Data are still very limited regarding the optimal management of transplant patients. Some centers (15-18) aggressively reduce the immunosuppression by withdrawing antimetabolites, discontinue or major reduction in calcineurin inhibitor (CNI) while keeping small doses of steroids. Attention must also be paid to the major interaction between protease inhibitors (lopinavir–ritonavir) and CNIs. Moreover, it seems necessary to reduce the dose of CNIs while using chloroquine or hydroxychloroquine. Frequent monitoring of CNI drug levels is recommended. However, Carbajo-Lozoya J et al. have shown that Coronavirus (CoV) replication depends on active immunophilin pathways. Tacrolimus can strongly inhibit the growth of the human coronaviruses SARS-CoV, HCoV-NL63, and HCoV-229E at low, non-cytotoxic concentrations in cell culture [19]. Based on these data, it makes sense to study the effect of keeping low doses of tacrolimus in Covid-19 infected SOT patients. In more advanced stages of Covid-19 (cytokine storm syndrome), treatment probably should focus on reducing uncontrolled inflammation by blocking IL-6, TNF alpha, or by removing cytokines by hemoperfusion.

To conclude, in populations at risk (elderly, associated comorbidities, immunosuppressed), when activation of the innate immune system fails to produce an adequate adaptive response (i.e., virus-specific CD8<sup>+</sup> T-cells), it seems that persistent self-induced inflammation can then cause mortality. Thus, mounting an early adaptive immune response may save lives.

## References:

1. World Health Organization. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>, 19 March 2020.
2. Fung SY, Yuen KS, Ye ZW, Chan CP, Jin DY. A tug-of-war between severe acute respiratory syndrome coronavirus 2 and host antiviral defence: lessons from other pathogenic viruses, *Emerging Microbes & Infections*. 2020; 9(1):558-570. Doi: 10.1080/22221751.2020.1736644.
3. Jiankun Z, Chandra C. Toll-Like Receptor Signaling Pathways—Therapeutic Opportunities. *Mediators Inflamm*. 2010 ; 781235. Doi :10.1155/2010/781235.
4. Zhou Y, He C, Wang L, Ge B. Post-translational regulation of antiviral innate signaling. *Eur. J. Immunol*. 2017; 47(9): 1414-1426. Doi:10.1002/eji.201746959
5. Teijaro JR, Walsh KB, Rice S, Rosen H, Oldstone MB. Mapping the innate signaling cascade essential for cytokine storm during influenza virus infection. *Proc Natl Acad Sci. USA*. 2014;111(10):3799–3804. Doi:10.1073/pnas.1400593111.
6. Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA*. Published online March 23, 2020. doi:10.1001/jama.2020.4683
7. Weiskopf D, Weinberger B, Grubeck-Loebenstien, B. The aging of the immune system. *Transplant Int*. 2009; 22:1041-1050. Doi:10.1111/j.1432-2277.2009.00927.x.
8. Velazquez-Salinas L, Verdugo-Rodriguez A, Rodriguez LL, Borca MV. The Role of Interleukin 6 During Viral Infections. *Front. Microbiol*. 2019; 10:1057. Doi: 10.3389/fmicb.2019.01057.
9. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir–Ritonavir in Adults Hospitalized with Severe Covid-19. *NEJM*. 2020 March 18. Doi: 10.1056/NEJMoa2001282.
10. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-labeled non-randomized clinical trial. *Int J Antimicrob Agents*. 2020 Mar 20 :105949. Doi: 10.1016/j.ijantimicag.2020.105949.
11. Devaux CA, Rolain JM, Colson P, Raoult D. New insights on the antiviral effects of chloroquine against coronavirus: what to expect for COVID-19? *Int J Antimicrob Agents*. 2020 Mar 11:105938. Doi: 10.1016/j.ijantimicag.2020.105938.
12. Zhou D, Dai SM, Tong Q. COVID-19: a recommendation to examine the effect of hydroxychloroquine in preventing infection and progression. *J Antimicrob Chemother*. 2020 Mar 20. Pii : dkaa114. Doi: 10.1093/jac/dkaa114.

- Accepted Article
13. Schultz MJ. Macrolide activities beyond their antimicrobial effects: macrolides in diffuse panbronchiolitis and cystic fibrosis. *J Antimicrob Chemother.* 2004 ;54(1):21–28. Doi: 10.1093/jac/dkh309.
  14. Xu X, Han M, Li T, et al. Effective treatment of severe COVID-19 patients with Tocilizumab. *chinaXiv:202003.00026v1.*
  15. Alberici F, Delbarba E, Manenti C, et al. Management of patients on dialysis and with kidney transplant during covid-19 coronavirus infection. <https://www.era-edta.org/en/wp-content/uploads/2020/03/COVID>.
  16. López-Oliva MO, González E, Miranda RJ, Jiménez C. Management of kidney transplant immunosuppression in positive coronavirus infection requiring hospital admission. <https://www.era-edta.org>.
  17. Zhu L, Xu X, Ma K, et al. Successful recovery of COVID-19 pneumonia in a renal transplant recipient with long-term immunosuppression. *Am J Transplant.* 2020 Mar 17. Doi: 10.1111/ajt.15869.
  18. Guillen E, Pineiro GJ, Revuelta I, et al. Case report of COVID-19 in a kidney transplant recipient: Does immunosuppression alter the clinical presentation? *Am J Transplant.* 2020 Mar 20. Doi: 10.1111/ajt.15874.
  19. Carbajo-Lozoya J, Müller MA, Kallies S, Thiel V, Drosten C, von Brunn A. Replication of human coronaviruses SARS-CoV, HCoV-NL63 and HCoV-229E is inhibited by the drug FK506. *Virus Res.* 2012 ;165(1):112-7. Doi : 10.1016/j.virusres.2012.02.002.